

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



B2

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68, C07H 21/02, 21/04, A61K 31/70, G01N 33/53	A1	(11) International Publication Number: WO 97/15685 (43) International Publication Date: 1 May 1997 (01.05.97)
(21) International Application Number: PCT/EP96/04671 (22) International Filing Date: 25 October 1996 (25.10.96) (30) Priority Data: 95116890.5 26 October 1995 (26.10.95) EP (34) Countries for which the regional or international application was filed: DE et al. (71)(72) Applicant and Inventor: WINNACKER, Ernst-Ludwig [DE/DE]; Dall'Armistrasse 41a, D-80638 München (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): WEISS, Stefan [DE/DE]; Am Schorn 28, D-82327 Tutzing (DE). FAMULOK, Michael [DE/DE]; Schmaedelstrasse 28, D-81245 München (DE). (74) Agent: VOSSIUS & PARTNER; P.O. Box 86 07 67, D-81634 München (DE).		(81) Designated States: AL, JP, LT, LV, RO, SI, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: NUCLEIC ACID MOLECULES CAPABLE OF DISTINGUISHING THE ISOFORMS PrP ^C AND PrP ^{Sc} OF PRION PROTEINS AND PROCESSES FOR THEIR PRODUCTION (57) Abstract The invention describes a process for the identification and isolation of nucleic acid molecules capable of distinguishing the isoforms PrP ^C and PrP ^{Sc} of prion proteins as well as nucleic acid molecules obtainable by this process. Furthermore, pharmaceutical compositions and diagnostic compositions are described which comprise nucleic acid molecules specifically binding prion protein isoforms as well as diagnostic methods using such molecules.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Nucleic acid molecules capable of distinguishing the isoforms PrP^C and PrP^{Sc} of prion proteins and processes for their production

The present invention relates to a process for the identification and isolation of nucleic acid molecules capable of distinguishing the isoforms PrP^C and PrP^{Sc} of prion proteins as well as to the nucleic acid molecules obtainable by this process. Furthermore, the invention relates to pharmaceutical and diagnostic compositions comprising said nucleic acid molecules.

Proteinaceous infectious particles called prions are thought to be the causative agent of transmissible spongiform encephalopathies (TSEs) such as Scrapie of sheep, bovine spongiform encephalopathy (BSE) of calf, transmissible mink encephalopathy (TME) of mink as well as Kuru, Gerstmann-Sträussler-Scheinker syndrome (GSS), Creutzfeldt-Jakob-Disease (CJD) and fatal familial insomnia (FFI) in the case of humans (Prusiner, 1982). The main component of prions associated in amyloid-like rods (Prusiner et al., 1983; 1984) or scrapie associated fibrils (SAF; Hope et al., 1986) was found to be the prion protein PrP²⁷⁻³⁰ (Prusiner et al., 1981; Prusiner et al., 1983), an N-terminal truncated, highly protease resistant version of the prion protein PrP^{Sc} (Oesch et al., 1994), which is also found to a minor extent in prion preparations (Prusiner et al. 1983). PrP²⁷⁻³⁰, which is devoid of 67 amino acids at the aminoterminal end, results from PrP^{Sc} by proteinase K digestions (Prusiner et al., 1984; Stahl et al., 1993) or by lysosomal protease digestion (Caughey et al., 1991). The distribution of PrP^{Sc} and PrP²⁷⁻³⁰ in prion preparations varies dependent from the absence or presence of proteases.

No specific nucleic acid could be detected so far in prion preparations (Kellings et al., 1992) suggesting that the prion is infectious and can replicate in the absence of any nucleic acid (Prusiner, 1982). According to the protein-only hypothesis (Prusiner, 1982) exogenous PrP^{Sc}/PrP²⁷⁻³⁰ could convert the ubiquitous cellular isoform PrP^C

to PrP^{Sc}/PrP27-30. It is assumed that chaperons may be involved in this process (Edenhofer et al., 1996). PrP^{Sc}/PrP27-30 could appear as a monomer (Prusiner, 1982) or as a nucleation or crystal seed consisting of a PrP^{Sc}/PrP27-30 oligomer (Lansbury and Caughey, 1995). PrP^C differs from PrP27-30 only with respect to its secondary structure: the α -helical and β -sheet contents of PrP^C are 42 % and 3 %, respectively (Pan et al., 1993). In contrast, the α -helical and β -sheet contents of PrP27-30 were proven to be 21 % and 54 %, respectively (Pan et al., 1993). These results indicate that the conversion of PrP^C to PrP^{Sc}/PrP27-30 is most likely concomitant with extreme alterations in the secondary structure of the prion protein.

Although a series of experiments employing knock-out mice which no longer express PrP^C suggest that cellular prion proteins could play a crucial role in a number of cellular processes (Collinge et al., 1994; Sakaguchi et al., 1996; Tobler et al., 1996), a precise physiological role of PrP^C remains speculative. It is, however, proven that PrP^C is necessary for the development of transmissible spongiform encephalopathies (Büeler et al., 1993; Brandner et al., 1996).

Translation of the mRNA from Scrapie infected Syrian golden hamster has led to a 254 amino acid protein including a 22 amino acid signal peptide at the NH₂-terminus and a 23 amino acid signal sequence at the carboxy terminus (Oesch et al., 1985; Basler et al., 1986). The mature protein PrP^C as well as the Scrapie isoform PrP^{Sc} contain amino acids 23 to 231. Only PrP^{Sc} can be processed to the proteinase K resistant isoform PrP27-30 (amino acids 90-231) consisting of 142 amino acids (Prusiner et al., 1984).

This property has been used to design a diagnostic assay for diseases in connection with prion proteins in which a probe is treated with proteinase K in order to degrade all PrP^C and then reacted with an antibody directed against prion proteins (Groschup et al., 1994). However, this assay has the disadvantage that sensitivity might be hampered by the fact that the proteinase K digestion of PrP^C is not complete, thereby leading to false positive results. Furthermore, the additional step of proteinase K digestion is time consuming. In order to be able to directly assay for the presence or absence of PrP^C and/or PrP^{Sc} one would need antibodies which could distinguish between these two isoforms.

However, so far attempts to provide antibodies that can distinguish between the cellular isoform PrP^c and the isoforms PrP^{Sc}, as well as the truncated version PrP27-30, have failed (Groschup et al., 1994 and ref. therein).

Thus, up to now it was not possible to distinguish the isoforms PrP^c and PrP^{Sc} of prion proteins by immunological or other means which would be the prerequisite for a simple and reliable method of diagnosing a transmissible spongiform encephalopathy.

Therefore, the technical problem underlying the present invention is to provide a process for the identification and isolation of molecules which are capable of distinguishing between the isoforms PrP^c and PrP^{Sc} or PrP27-30 of prion proteins and which are useful tools for diagnosis and therapy of transmissible spongiform encephalopathies.

The solution to said technical problem is achieved by the provision of the embodiments characterized by the patent claims.

Thus, the present invention relates to a process for the identification and isolation of nucleic acid molecules which are capable of distinguishing between the isoforms PrP^c and PrP^{Sc} or PrP27-30 of prion proteins associated with transmissible spongiform encephalopathies comprising the steps of

- (i) incubating a prion protein isoform or peptide fragment or derivative of this prion protein isoform with a pool of nucleic acid molecules comprising different sequences;
- (ii) selecting and isolating those nucleic acid molecules which are capable of binding to said prion protein isoform or fragment or derivative thereof;
- (iii) optionally, amplifying the isolated nucleic acid molecules and repeating steps (i) and (ii); and
- (iv) determining the binding specificity of the isolated nucleic acid molecules for the PrP^c and PrP^{Sc} or PrP27-30 isoforms of prion proteins.

The process according to the invention is based on a method called "in vitro selection". This method allows for the identification of nucleic acid molecules (RNA, modified RNA, ssDNA or dsDNA) which bind with high affinity to a defined molecular target from a large randomized population of nucleic acid molecules (Tuerk and Gold, 1990; Famulok and Szostak, 1992). Using this method it has been possible to isolate nucleic acids specifically recognizing a variety of protein targets including HIV-1 reverse transcriptase (Tuerk et al., 1992), HIV-1 Integrase (Allen et al., 1995), human α -thrombin (Kubik et al., 1994) and *Drosophila* sex-lethal protein (Sakashita and Sakamoto, 1994). However, up to now it has not been possible to provide by this method nucleic acid molecules being capable of distinguishing the two isoforms of prion proteins, PrP^c and PrP^{Sc}. In the scope of the present invention the term PrP^c comprises the cellular isoform of the prion protein as well as fragments and derivatives thereof irrespective of the source organism. The term PrP^{Sc} comprises the isoform of the prion protein associated with various transmissible spongiform encephalopathies. This term also comprises fragments of this prion protein isoform such as the truncated version of the isoform PrP^{Sc}, the prion protein PrP27-30, which is the main component of prions. In particular, this term also includes PrP^{Sc} proteins of the various Scrapie strains including those adapted to hamster, mouse or other vertebrates. Also included are derivatives of the prion protein isoform PrP^{Sc}.

The term derivatives includes chemically modified versions of the prion protein isoforms PrP^c and PrP^{Sc} as well as mutants of these proteins, namely proteins which differ from the naturally occurring prion protein isoforms at one or more positions in the amino acid sequence, as well as proteins that show deletions or insertions in comparison to the naturally occurring prion protein isoforms. Such mutants can be produced by recombinant DNA technology or can be naturally occurring mutants. The term derivatives also embraces proteins which contain modified amino acids or which are modified by glycosylation, phosphorylation and the like.

According to the invention it is possible to use as nucleic acid molecules single or double stranded nucleic acid molecules, such as RNA, modified RNA, single stranded DNA or double stranded DNA.

A pool of nucleic acid molecules, which constitutes the starting material from which nucleic acid molecules are selected which specifically bind to one of the isoforms of the prion protein, is defined as a mixture of nucleic acid molecules of different sequences. This pool can be any mixture of nucleic acid molecules, preferably a pool of randomized molecules. Preferably the nucleic acid molecules of the pool are chemically synthesized or produced by in vitro transcription.

In the case of RNA molecules the RNA pool which is screened for molecules specifically binding to one of the isoforms of a prion protein is preferably the RNA pool M111.1 described in Famulok (1994). This pool consists of RNA molecules of 111 nucleotides randomized at 74 positions and results from the transcription of corresponding DNA sequences. The pool M111.1 contains RNA molecules with approximately 1×10^{15} different sequences.

The process according to the invention can be used to identify and isolate nucleic acid molecules which can distinguish between the two isoforms of prion proteins, PrP^c and PrP^{Sc}, associated with a transmissible spongiform encephalopathy such as Scrapie of sheep, bovine spongiform encephalopathy (BSE) of calf, transmissible mink encephalopathy (TME) of mink, Kuru, Gerstmann-Sträussler-Scheinker Syndrome (GSS), fatal familial insomnia (FFI), Creutzfeldt-Jakob Disease (CJD) in the case of humans, chronic wasting disease (CWD) of mule, deer and elk or feline spongiform encephalopathy (FSE) of cats. Transmissible spongiform encephalopathies are also known from nyala, gemsbok, arabian oryx, greater kudu, eland, ankole, moufflon, puma, cheetah, scimitar horned oryx, ocelot and tiger.

The step of incubating the pool of nucleic acid molecules with a prion protein can be carried out in different ways.

In one preferred embodiment of the invention the protein is immobilized, for example, on a matrix such as a gel or a resin for chromatography. The immobilization can be achieved by means known to the person skilled in the art. For example, the protein can be covalently linked to a matrix or can be bound to it by a specific interaction between a group present on the matrix and a domain of the protein specifically

recognizing this group. Such a domain can be fused to a prion protein by recombinant DNA technology as will be discussed below.

If the prion protein is immobilized, nucleic acid molecules which do not bind to the prion protein can be removed after incubation by washing with an appropriate buffer. Subsequently the nucleic acid molecules binding to the prion protein can be eluted from the immobilized protein, for example by 8M urea, and further purified, for example, by phenol extraction and precipitation.

In another preferred embodiment the prion protein is in solution. In this case the nucleic acid molecules binding to the prion protein can be isolated, for example, by carrying out a gel retardation assay and isolating the protein/nucleic acid complex. Subsequently the nucleic acid molecules can be isolated from the complex and further purified by known methods.

According to the invention it is possible to amplify the nucleic acid molecules obtained by steps (i) and (ii), for example by in vitro transcription, reverse transcription or polymerase chain reaction or a combination of these techniques, and to repeat steps (i) and (ii). This leads to a further selection and amplification of nucleic acid molecules which bind specifically to the used prion protein.

If several cycles of steps (i) to (iii) of the process are performed, it is possible to use in one or more cycles an immobilized protein and in one or more cycles a protein in solution. A cycle in which a protein in solution is used permits the elimination of nucleic acid molecules binding to the matrix on which the immobilized protein is fixed.

The prion protein used in the process can be any of the known prion protein isoforms or a fragment or derivative of such a protein.

In a preferred embodiment the prion protein is the isoform PrP^{Sc} present in the prion. In a specifically preferred embodiment the N-terminally truncated version of PrP^{Sc}, PrP27-30, is used. In this context PrP^{Sc} and PrP 27-30 refer to any of these isoforms which can be found in an organism affected with a transmissible spongiform encephalopathy.

In a further preferred embodiment the prion protein used in the process is the cellular isoform PrP^c, most preferably the processed form PrP^c23-231 which comprises amino acids 23 to 231 of PrP^c.

In another preferred embodiment the prion protein used in the process is a recombinant protein. This means that the protein is produced by recombinant DNA technology, namely by expression from a cloned DNA sequence.

More preferably, the prion protein is part of a fusion protein. Such a fusion protein can comprise beside the prion protein a protein or protein domain which confers to the fusion protein a specific binding capacity. For example, such a domain may be an oligohistidine (Le-Grice et al., 1990), Calmoduline binding peptide (CBP) (Carr et al., 1991), S-peptide (ribonuclease A) (Kim and Raines, 1993), FLAG (Kawase et al., 1995), green-fluorescent protein (GFP) (Hampton et al., 1996), BTag (Wang et al., 1996), or maltose-binding protein (MBP) (Aitken et al., 1994, Richards and Wyckoff, 1971). Proteins comprising such a domain can be immobilized for example, on IMAC-Ni²⁺, Calmodulin, S-protein 104 aa (Kim and Raines, 1993), anti-FLAG-antibodies, anti-GFP-antibodies, BTag-antibodies or maltose. Elution can then be achieved by a method well-known in the art. In a preferred embodiment the prion protein is fused to glutathione-S-transferase. Such a fusion protein possesses a high affinity for glutathione and can thus be immobilized on a matrix comprising glutathione, such as glutathione-sepharose.

In the last step of the process according to the invention the isolated nucleic acid molecules are tested for their binding to the different isoforms, PrP^c and PrP^{Sc}, of a prion protein. Those nucleic acid molecules are selected which specifically bind to only one of the isoforms.

Thus, the process according to the invention allows the identification and isolation of nucleic acid molecules which specifically bind to one of the isoforms of a prion protein or a fragment or derivative thereof and thereby allow the distinguishing of the different isoforms. These nucleic acid molecules therefore show an unexpected high

specificity, which is even higher than the specificity of poly- or monoclonal antibodies which cannot distinguish between the isoforms of prion proteins.

The process of the invention has been successfully carried out to isolate RNA molecules which can distinguish between the isoforms PrP^c23-231 and PrP27-30 from Syrian Golden Hamster. In this case the isoforms were recombinant proteins fused to glutathione-S-transferase (GST::PrP^c23-231 and GST::rPrP27-30). The recombinant rPrP27-30 protein is identical in sequence to the natural PrP27-30 protein but reveals in contrast to the natural isoform proteinase K sensitivity.

Furthermore, the present invention relates to nucleic acid molecules obtainable by a process according to the invention, namely to RNA, single stranded DNA or double stranded DNA molecules which bind to one of the isoforms of a prion protein. These include nucleic acid molecules which specifically bind to the cellular isoform PrP^c, namely to the processed form PrP^c23-231, or specifically to the isoform PrP^{Sc}, namely to the truncated version PrP27-30, or specifically to derivatives of these proteins.

In a preferred embodiment the nucleic acid molecules of the invention comprise four stretches of three consecutive guanosine residues separated by single stranded regions between four and seven nucleotides long. More preferably, the nucleic acid molecules comprise a nucleotide sequence as depicted in SEQ ID NO: 15, SEQ ID NO: 16 or SEQ ID NO: 17.

In another preferred embodiment, the region comprising the four guanosine stretches is flanked by two variable regions of predominantly Watson-Crick covariation. In particular, the nucleic acid molecules preferably comprise a nucleotide sequence as depicted in any one of SEQ ID NO: 1 to 13 and more preferably a nucleotide sequence as depicted in SEQ ID NO: 18.

In a preferred embodiment the nucleic acid molecules according to the invention are further modified at one or more positions in order to increase their stability and/or to alter their biochemical and/or biophysical properties.

The present invention also relates to pharmaceutical compositions comprising nucleic acid molecules according to the invention. Such compositions can optionally comprise pharmaceutically acceptable carriers.

These compositions may be useful for the therapy of transmissible spongiform encephalopathies such as those listed above. It may be possible, for example, to suppress the conversion of the isoform PrP^c into the prion associated isoform PrP^{Sc} by applying nucleic acid molecules which specifically bind to PrP^c .

Furthermore, the present invention relates to diagnostic compositions comprising nucleic acid molecules according to the invention. Such compositions may contain additives commonly used for diagnostic purposes. The nucleic acid molecules and the diagnostic compositions according to the invention can be used in methods for the diagnosis of transmissible spongiform encephalopathies. Such a method comprises, for example, the incubation of a probe taken from a body with at least one kind of nucleic acid molecules according to the invention and the subsequent determination of the interaction of the nucleic acid molecules with the isoforms PrP^c and PrP^{Sc} of a prion protein.

Since during the occurrence of a transmissible spongiform encephalopathy the amount of the isoform PrP^{Sc} increases and the total amount of the cellular isoform PrP^c decreases, it is in principle possible to use for diagnosis nucleic acid molecules which bind to one or the other of the two isoforms.

On the one hand, it is possible to use at least one kind of nucleic acid molecule according to the invention in order to quantitatively determine the amount of at least one isoform of a prion protein in a probe.

On the other hand, it is possible to use nucleic acid molecules which specifically bind the PrP^c isoform in combination with nucleic acid molecules which specifically bind the PrP^{Sc} isoform in order to determine the absolute and/or relative amount of the isoforms in a probe.

In a preferred embodiment the probe may be obtained from various organs, preferably from tissue, for example, from brain, tonsils, ileum, cortex, dura mater, Purkinje cells, lymphnodes, nerve cells, spleen, muscle cells, placenta, pancreas,

eyes, backbone marrow or peyer'sche plaques, for example in the form of thin sections. Alternatively the probe may be obtained from a body fluid, preferably from blood, cerebrospinal fluid, milk or semen.

In the case that brain is used as a probe, diagnosis is in most cases performed post mortem. Exceptionally, brain biopsies can be performed on the alive organism. The brain can originate from any organism that might be afflicted with a transmissible spongiform encephalopathy, such as sheep, calf, mice, cats, hamster, mule, deer, elk or humans or from other organisms which may be afflicted by a TSE as mentioned above. The brain should originate from organisms which are PrP^{0/0} (knock-out), PrP^{Sc} (infected) and PrP^C (wild-type) or of unknown PrP-status.

In the case that blood, milk, cerebrospinal fluid, semen or tissue from other organs as mentioned above is used as a probe, diagnosis is possible for living individuals.

Furthermore, the nucleic acid molecules according to the invention can be used to identify three dimensional structures which are necessary for the specific binding of a prion protein isoform. With the help of this information other chemical compounds can be isolated or synthesized which can specifically bind prion protein isoforms. Thus, the present invention also relates to chemical compounds other than nucleic acid molecules which are based on the information derived from a three dimensional structure of a nucleic acid molecule according to the invention, selected from the group consisting of inorganic or organic compounds, preferably sugars, amino acids, proteins or carbohydrates.

Figure 1A: Illustrates schematically the method for in vitro selection of RNA molecules specifically binding to the immobilized fusion protein GST::rPrP^C (GST:: rPrP23-231) (GST = glutathione-S-transferase; PCR = Polymerase chain reaction).

In the following, (r)PrP^C stands for rPrP23-231. Furthermore, rPrP27-30 stands for rPrP90-231 (Syrian Golden Hamster).

- Figure 1B:** Illustrates schematically a further step in the *in vitro* selection of RNA molecules specifically binding to GST::rPrP^c using GST::rPrP^c in solution and a gel retardation assay.
- Figure 2:** Schematically illustrates the construction by *in vitro* transcription of the randomized RNA pool M111.1 (Famulok, 1994) (Ntes = nucleotides).
- Figure 3:** Shows the percentage of RNA binding to immobilized GST::rPrP^c23-231 after each cycle of the process described in Example 1. Radioactivity associated with GST::rPrP^c beads after removal of the supernatant was set to 100 %. Radioactivity retained after 4 washing steps represents the percentage of RNA binding.
- Figure 4A:** Shows the binding of selected RNAs and unselected RNAs to GST, GST::rPrP^c and GST:rPrP27-30. 5' labeled RNA was incubated in the presence of the proteins, filtered over BA85 nitrocellulose on a millipore slot blot apparatus. Retained radioactivity was quantified by Cerenkov counting.
- Figure 4B:** Shows that the *in vitro* selected RNA molecules of Example 1 distinguish between PrP^c and rPrP27-30 from the Syrian Golden Hamster. Gel a: 5' labeled RNA molecules after 9 cycles were incubated in the presence of GST, GST::PrP^c and GST::rPrP27-30, and analyzed on 0.7 % non-denaturing agarose gels. Gels were fixed by 5 % TCA, dried and subjected to autoradiography. The GST::rPrP^c/RNA complex from the 9th cycle was excised from the gel, the RNA extracted, reverse transcribed, PCR amplified and *in vitro* transcribed (see Figure 1B). This procedure was repeated twice for cycle 10 and 11. Gel b: The 5' labeled RNA - after 11 cycles - was again incubated in the presence of GST, GST::rPrP^c and GST::rPrP27-30 and analyzed as described above.

Figure 5: Sequences of selected RNA aptamers directed against rPrP^C23-231 fused to GST from hamster by *in vitro* selection. The aptamers belong to several groups of molecules. RNA aptamers of group (A) (motif I) and (B) (motif II) can harbor G-quartet motifs and distinguish between rPrP23-231 (rPrP^C) and rPrP90-231 (rPrP27-30). RNA aptamers of group (C) (motif III) could also have G-quartet motifs but interact with rPrP23-231 (rPrP^C) and rPrP90-231 (rPrP27-30). (D) Aptamers with unique G-quartets (5 out of 6 aptamers shown). Aptamers of group (E) lack any G-quartet motif and bind to GST (one out of 6 aptamers shown).

Figure 6: RNA aptamers motif I and II distinguish the recombinant prion protein isoforms rPrP23-231 (rPrP^C) and rPrP90-231 (rPrP27-30) from hamster and calf. (A) 4 pMols of labeled RNA Ap1 (motif I; lanes 1-3) were incubated in the presence of 40 pMols each of recombinant GST::rPrP23-231 (rPrP^C) (lane 2) and GST::rPrP90-231 (rPrP27-30) from Syrian golden hamster (lane 3). (B) 4 pMols of labeled RNA Ap1 (motif I; lanes 1-3) were incubated in the presence of 40 pMols each of recombinant GST::bov-rPrP25-242 (rPrP^C) (lane 2) and GST::bov-rPrP93-242 (rPrP27-30 + 1 octarepeat) from calf (lane 3). (C) 4 pMol of labeled RNA motif II (lanes 1-3) were incubated in the presence of GST::rPrP23-231 (rPrP^C) (lane 2) and GST::rPrP90-231 (rPrP27-30) (lane 3) from hamster. Reaction assays were analyzed on 0.7 % agarose gels. (D) 4 pMols of labeled RNA Ap2 (motif II; lanes 1-3) were incubated in the presence of 40 pMols each of recombinant GST::bov-rPrP25-242 (rPrP^C) (lane 2) and GST::bov-rPrP93-242 (rPrP27-30 + 1 octarepeat) from calf (lane 3). The additional bovine octarepeat extends from aa 93 to 101.

Figure 7: Mapping of the RNA aptamer - PrP interaction site of hamster and calf. (A) 4 pMols of labeled RNA aptamer motif I (lanes 1-9) and (B) 4 pMols of labelled RNA aptamer motif II (lanes 1-9) were incubated in the

presence of 40 pMol each of GST::rPrP23-231 (rPrP^C) (lanes 8), GST::rPrP90-231 (rPrP27-30) (lanes 9) from hamster and 20 pMol each of GST::P23-52 (lanes 2), GST::P55-93 (lanes 3), GST::P90-109 (lanes 4), GST::P129-175 (lanes 5), GST::P218-231 (lanes 6) and GST::P180-210 (lanes 7). Reaction assays were analyzed on 0.7 % agarose gels. (Top) Schematic presentation of the hamster PrP region. Hatched box, PrP region interacting with the aptamers. Void boxes, PrP region not interacting with the aptamers. (C) 4 pMols of labelled RNA aptamer motif II (lanes 1-4) were incubated in the presence of 40 pMol each of bovine GST::bovP₂₅₋₉₂ (lane 2), GST::bovP₉₃₋₁₂₀ (lane 3) and 20 pMol each of bovine GST::bov-rPrP93-242 (rPrP27-30 + 1 octarepeat; lane 1) and hamster GST::P23-89 (lane 4).

The Examples illustrate the invention.

Example 1:

***In vitro* Selection of RNA molecules specifically binding GST::rPrP^C23-231**

An *in vitro* selection procedure (schematically outlined in Figure 1A and B) was carried out using recombinant PrP23-231 (rPrP^C) from the Syrian Golden Hamster fused to GST (Weiss et al., 1995) and RNA pool M 111.1 (Famulok, 1994).

Cycles 1-9: 5' [γ -³²P]-ATP labeled (1.Cycle) or [α -³²P]-UTP labeled (Cycles 2-10) RNA M111.1 (Famulok, 1994) (6,8 nMol (first), 1.82 nMol (2nd), 914 pMol (3rd), 665 pMol (4th), 2.07 nMol (5th), 831 pMol (6th), 2.7 nMol (7th), 1.94 nMol (8th and 9th cycle) was incubated in the presence of immobilized GST (185 pMol) synthesized in the Baculovirus system (Weiss et al., 1995) in binding buffer comprised of 8 mM Na₂HPO₄, 0.87 mM KH₂PO₄, 136 mM NaCl, 112.6 mM KCl, 2 mM DTT and 2 mM MgCl₂ (Fig. 1A). Incubation was done at 37°C in an overhead incubator.

Cycle 1-7: After 60 min. beads were collected for 10 min. at 700 g.

Cycle 8-9: Incubation with immobilized GST was done for 30 min. as described above. Subsequently the beads were removed by centrifugation and the supernatant incubated with freshly immobilized GST for another 30 minutes.

Cycle 1-9: The supernatant from the preselection(s) was incubated with immobilized GST::rPrP^c23-231 (53 pMol) synthesized in the Baculovirus system (Weiss et al., 1995) as described above. After 60 minutes the beads were washed four times with binding buffer and the RNA eluted in the presence of 8 M urea in 100 mM sodium citrate pH 8.0 and 3 mM EDTA. The RNA was phenol (pH 5.0)/chloroform extracted and EtOH precipitated in the presence of 2 M NH₄-acetate (Fig. 1A).

Cycle 9 to 11: Selected RNA (40 pMol in the 9th; 4 pMol each in the 10th and 11th cycle) was 5' labeled and incubated with soluble GST::rPrP^c23-231 (Weiss et al., 1995; 140 pMol in the 9th cycle, 40 pMol in the 10th and 11th cycle) for 60 min. at 37°C in binding buffer and analyzed by an gel retardation assay on an 0.7 % native agarose gel (Fig. 1B) as described (Weiss et al., 1992). Following electrophoresis the RNA/GST::PrP^c23-231 complex was excised and extracted by employing an Qiaex extraction kit (Qiagen).

Cycle 1-11: 50 % each of the extracted RNA was subjected to a reverse transcription reaction according to the Superscript reverse transcriptase kit (Gibco, BRL). 50 % of the resulting cDNA was amplified by PCR according to Saiki et al., 1988 using the primers shown in Figure 2B. 50 % of the amplified cDNA was *in vitro* transcribed as described (Weiss et al., 1992).

Nitrocellulose binding assay: 4 pMol 5' labeled (Sambrook et al., 1989) RNA was incubated in the presence of 0 to 500 nM of protein in the presence of binding buffer for 60 min. at 37°C. The incubation mixture was filtered over a BA85 nitrocellulose membrane in a Millipore slot blot apparatus, the filter washed with 4 ml of incubation buffer, excised and measured by Cerenkov Counting.

Gel retardation assay: RNA and protein were incubated as described above and the reaction mixture loaded on a 0.7 % native agarose gel as described (Weiss et al., 1992). Following electrophoresis, the gel was fixed by 5 % TCA, dried and subjected to autoradiography.

RNA Pool M111.1: The RNA pool was prepared by *in vitro* transcription from a DNA pool (138 bases) as described (Famulok, 1994). In brief, M111.1 reveals a randomized sequence of 74, a base permutation of $474 = 3.56 \times 10^{44}$ molecules, a

molecular weight of 36630 Dalton. Synthesis yielded 175 µg (4.76 nMol) RNA; that is 6×10^{23} (Avogadro) $\times 4.76 \times 10^{-9} = 2.86 \times 10^{15}$ molecules. 36 % of the synthesized ssDNA pool are extendible by PCR which resulted in a pool with approximately 1.03×10^{15} different sequences and a complexity of 1.03×10^{15} , which is equivalent to one pool copy (i.e. each individual RNA molecule is represented one fold in the pool). In particular, the technical features of RNA pool M111.1 are the following:

Nucleotide sequence of fixed region 1

5' CCGAATTCTAATACGACTCACTATAGGAGCTCAGC
CTTCACTGC (SEQ ID NO: 19)

Nucleotide sequence of fixed region 2

5' GTGGATCCGACCGTGGTGCC (SEQ ID NO: 20)

randomized sequence = 74 nucleotides

base permutation $4^{74} = 3.56 \times 10^{44}$;

MW = 36630; 1nM = 36.63 µg;

175µg (4.76 nMol) were synthesized; that is $6 \times 10^{23} \times 4.76 \times 10^{-9} = 2.86 \times 10^{15}$ molecules

36 % extendable

pool complexity = 1.03×10^{15} molecules

= 1 pool copy i.e. each individual RNA molecule is represented one fold in the pool

A schematic view of RNA pool M111.1 is shown in Fig. 2. The nucleotide sequence 5'-CCGAATTCTAATACGACTCACTATA (nucleotides 1 to 25 of SEQ. ID NO: 19) of the fixed region 1 only belongs to the DNA pool since it is not transcribed.

After 9 rounds of selection 7.2 % of the selected RNA bound to GST::rPrP^c23-231 immobilized on glutathione-sepharose 4B (Table 1 and Fig. 3).

The percentage of RNA-binding was determined as follows: Radioactivity associated with GST::rPrP^c beads after removal of supernatant was set to 100 %. Radioactivity retained after four washing steps represents the percentage of the RNA binding.

Table 1:

RNA protein ratios and % binding of RNA to protein dependent on the selection cycle

Cycle	RNA:GST (molar ratio)	RNA: GST::rPrP ^c (molar ratio)	% binding* (RNA to GST::rPrP ^c)
1	36:1	83:1	0
2	10:1	22:1	0
3	5:1	11:1	0
4	3.5:1	8:1	1
5	10:1	24:1	1.25
6	4.5:1	10:1	2.5
7	14:1	33:1	2.75
8	10:1/7:1	18:1	7.2
9	10:1/7:1	18:1	7.2

* The percentage of RNA binding was measured as described in the legend to Figure 3.

A binding assay employing soluble GST, GST::PrP^c23-231 and GST::rPrP27-30 revealed that 2 % of the enriched RNA from cycle 9 bound to GST::PrP^c23-231 whereas only 1.1 % bound to GST::rPrP27-30 and GST (Fig. 4 A). This result indicates that ~5 % of the RNA bound to the matrix, i.e. glutathione-sepharose 4B. After 6 rounds of selection only 1 % of the RNA bound to immobilized GST::PrP^c23-231 and 0.7 % to GST (Fig. 4A).

A gel retardation assay with RNA isolated after 9 cycles of selection confirms that about 2 % of the RNA bound to GST::PrP^c23-231 at a molar ratio of 10:1 (protein:RNA) (Fig. 4B, panel a, lane 6), whereas no binding occurs under identical conditions in the case of GST::rPrP27-30 (lane 9) and GST (lan 3).

To enrich RNA specifically binding to GST::PrP^c23-231 and to remove RNA molecules binding to the matrix, we excised the RNA/GST::PrP^c23-231 complex, extracted and amplified the RNA (Fig. 1B) and subjected it to two further gel retardation assays. As demonstrated in Fig. 4B after a total of 11 cycles (panel b) we isolated an RNA which binds specifically to GST::PrP^c23-231 at a 10 fold molar excess of protein over RNA (Figure 4B, panel b, lane 6). No binding occurs in the presence of GST::rPrP27-30 (lane 9) and GST (lane 3). A more detailed binding analysis revealed that binding of RNA to GST::PrP^c23-231 occurs at a molar ratio (RNA:protein) between 1:1 and 5:1. These findings demonstrate the selection of an RNA which can distinguish between GST::PrP^c23-231 and GST::rPrP27-30.

These results demonstrate that it is possible to isolate an RNA aptamer (aptus = to fit) by *in vitro* selection which binds specifically to the cellular prion protein isoform PrP^c23-231 fused to GST. This RNA does not bind to the recombinant prion protein rPrP27-30 fused to GST and not to GST. Therefore a RNA was selected which can distinguish between PrP^c23-231 and rPrP27-30. Recombinant PrP27-30 share the same amino acid sequence compared to natural PrP27-30 present in Scrapie prion preparations (Prusiner et al., 1984) but reveals in contrast to the natural isoform proteinase K sensitivity.

The RNA aptamer able to distinguish between PrP^c23-231 and rPrP27-30 overcomes the problem that it is not possible to produce poly- and monoclonal PrP antibodies which recognize specifically only one PrP isoform (Groschup et al., 1984 and ref. therein) and provides a suitable tool for a reliable diagnostic of transmissible spongiform encephalopathy.

Example 2

Determination of sequences of the identified RNA molecules

In order to determine the sequences of RNA molecules identified after 11 cycles of amplification and selection, these RNA molecules were reverse transcribed into cDNA and amplified by PCR (Sambrook et al., 1989). The obtained cDNA was

restricted with EcoRI and BamHI, subcloned into pGEM-3-Zf(-) and the sequence of 20 different cDNA clones pGEM-Ap 1 to 20 determined according to Sanger et al. (1977). Sequences of 14 RNA molecules identified are depicted in Figure 5 (SEQ ID NO: 1 to 14). The obtained monoclonal RNAs revealed sequences which may contain G-quartet motifs. Three classes of G-quartet motifs (Table II; Fig. 5A, B, C) could be identified with more than one monoclonal RNA. 30 % of the sequenced DNA molecules encode for unique RNA molecules which may also contain G-quartets (Fig. 5D), 30 % of the selected RNA aptamers did not contain any G-quartet motif (Fig. 5E).

Table II. Distribution of selected RNA aptamers.

	Motif I	Motif II	Motif III	G-quartet motif (unique)	no G-quartet motif
% of clones sequenced	10	15	15	30	30

A detailed analysis of the 20 sequenced clones revealed that 70 % of the clones contained four sets of three highly conserved consecutive guanosine residues, separated by single stranded regions between four and seven nucleotides long. These guanosine rich consensus motifs are flanked by two variable regions of predominantly Watson-Crick variation (see Figure 6). The primary sequence of the molecules comprising the four sets of guanosine stretches strongly suggests that their secondary structure contains a three layered G-tetrad motif (see Figure 6). In 40 % of the selected RNAs three classes of aptamer motifs were identified based on their relationship within the three single stranded loop regions (see Figure 6). While

individual members of each identified class were identical in the putative G-tetrad and loop regions. they showed significant covariation in the Watson-Crick helix.

Such G-tetrad motifs had already been identified in several other in-vitro selected nucleic acid molecules (see e.g. Bock et al., 1992; Wang et al., 1993; Macaya et al., 1993; Lauhon and Szostak, 1995; Huizenga and Szostak, 1995; Harada and Frankei, 1995) and appear to represent an important feature in nucleic acid molecules which bind to a ligand with high specificity. Furthermore, G-quartets have been suggested for telomeric DNA sequences in species such as *Tetrahymena* (Sundquist and Klug, 1989; Williamson et al., 1989; for review: Williamson, 1993). Guanine rich sequences which could form G-quartets were found in immunoglobulin switch regions, gene promoters and in chromosomal telomers which are thought to bring the four homologous chromatids together during meiosis and prevent the DNA from degradation (Sen and Gilbert, 1988). G-quartets have also been discussed to play a role in the dimerization process of retroviral genomic RNA (Weiss et al., 1993), a prerequisite for the generation of infectious virions. G-tetrads are held together by Hoogsteen base pairing through hydrogen bonds between nitrogens or oxygens and hydrogens (Sen and Gilbert, 1988). The RNA aptamers selected against the prion protein could contain three G-quartets stacked upon each other (Fig. 6) to form two eight-coordinate chelation cages. Alkali-metal ions such as potassium located within the axial channel are able to complex four oxygens of the upper and four oxygens of the lower G-quartet. Because of the very compact structure G-tetrads are very stable and unusual RNase resistant.

Example 3

Monoclonal RNA aptamers harboring G-quartet motif I and II bind specifically to rPrP23-231 (rPrP^C) from hamster and rPrP25-242 (rPrP^C) from calf

Monoclonal RNA aptamers representing motif I (Ap1; Figure 6 A, B) and II (Ap2; Figure 6C, D) interact specifically with rPrP23-231 (rPrP^C) from Syrian golden hamster (Figure 6 A, C; lanes 2) and rPrP25-242 (rPrP^C) from cattle (Figure 6 B, D; lanes 2) both fused to GST. Prion proteins from hamster and cattle reveal a sequence homology of 88%. Bovine PrP was synthesized in insect cells infected with a recombinant baculovirus containing bovine prn-p cDNA (Yoshimoto et al., 1992). Binding of the RNA aptamers to bovine PrP was investigated to prove whether the aptamers are suitable for the development of a BSE diagnostic tool. Both aptamers do not bind to the recombinant GST fused prion proteins rPrP90-231 (rPrP27-30) from hamster (Figure 6, A, C; lanes 3) and rPrP93-242 (rPrP27-30 + 1 octarepeat, aa 93-101) from cattle (Figure 6, B, D; lanes 3) demonstrating that the molecules distinguish between rPrP^C and rPrP27-30 from both species. RNA aptamers of group III interact with rPrP^C and show weak interaction with rPrP27-30. Some of the selected RNA aptamers lacking any G-quartet motif interact with GST (data not shown).

Furthermore, an aptamer was constructed consisting of 60 nucleotides on the basis of an aptamer comprising motif I as shown in Figure 5, which, however, lacked the primer binding sites as well as 14 nucleotides of the randomized region. This 60-mer exactly corresponds to a part of one of the aptamers displaying the motif I as depicted in Figure 6A/B and displayed the same binding characteristics as the full-length aptamer. The sequence of this 60mer is depicted in SEQ ID NO: 18. This molecule was isolated by the following procedure: By using two appropriate primers,

DNA containing the 60 nucleotides encoding the RNA motif I (Figure 5A, #II; SEQ ID NO: 18) flanked by a T7 promoter was amplified by PCR (Sambrook et al., 1989) under the following conditions: 1 min 94°C, 1 min 52°C and 2 min 72°C for 25 cycles. The amplified cDNA product of 80 nucleotides was subjected to an in vitro transcription reaction (Sambrook et al., 1989) with T7 RNA polymerase leading to the RNA aptamer consisting of 60 nucleotides (motif I, Figure 6A, B). The RNA was gel purified before the use in gel retardation assays.

This 60mer was also used for the determination of the equilibrium binding constants. For this purpose 4 pMol of 5' γ -³²P-ATP labelled and α -³²P-UTP labelled RNA aptamer motif I (SEQ ID NO: 18) was incubated in the presence of 0, 4, 20, 28, 40, 60, 80, 108 pMol of GST::rPrP23-231 for 60 min at 37°C under assay conditions as described above. RNA/protein complexes have been analyzed by an gel retardation assay (Weiss et al., 1992). Gel was fixed by 5% TCA, dried and subjected to autoradiography for 12 hours. Intensities of the signals have been determined by phosphoimaging (ImageQuant™, Strom 860, Molecular Dynamics).

For the calculation of the equilibrium binding constant a bimolecular reaction between the RNA and the protein was assumed. The concentration [c] of the RNA/protein complex at equilibrium can be determined from the amount of radioactivity in the shifted position and the known specific activity of the RNA. For the calculation of the equilibrium binding constant (K_D) the following formula was used (Meisterernst et al., 1988; Schellenberger et al., 1989):

$$K_D = \left[\frac{R_o}{[PR]_{eq}} - 1 \right] \times [P]_{eq}$$

$$R_o = R_{eq} + [RP]_{eq};$$

$$P_{eq} = P_o - [RP]_{eq};$$

$$R = \text{RNA aptamer motif I (60mer)}$$

P = GST::rPrP23-231 (GST::rPrP^C);

The following equilibrium binding constant (K_D) for the complex of the RNA aptamer (60mer) and GST::rPrP23-231 (GST::PrP^C) was preliminary calculated as $K_D = 8 \times 10^{-7}$ M. Applying other models not basing on bimolecular binding reactions K_D values $< 8 \times 10^{-7}$ M are expected.

Example 4

Mapping of the hamster and bovine PrP/aptamer binding site

To map the interaction site of Syrian golden hamster PrP^C to RNA aptamers 1 and 2, we employed a series of recombinant prion peptides (Weiss et al., 1995; Figure 7, A, B). Only peptide P23-52 interact with RNA aptamers Ap1 and 2 (Figure 7, A, B), demonstrating that the amino terminal residues aa23 to aa52 of the Syrian golden hamster prion protein are sufficient for the recognition by both aptamers. Binding of the aptamers to rPrP27-30 failed because this molecule lacks the amino terminal 67 amino acid residues. Prion peptides P25-92 and P93-120 (Fig. 7C, lane 3) from bovine PrP have been synthesized to map the interaction site of bovine PrP^C to RNA aptamer motif II. Only P25-92 (Fig. 7C, lane 2) did bind to RNA aptamer motif II (Ap2) demonstrating that it is the amino terminus of the bovine prion protein which is recognized by the aptamer. Hamster peptide P23-89 (Fig. 7C, lane 4) did also interact with aptamer Ap2 confirming the interaction of the amino terminus of the hamster prion protein with aptamer motif II.

Cited References

Aitken, R., Gilchrist, J. and Sinclair, M.C.: Vectors to facilitate the creation of translational fusions to the maltose-binding protein of *Escherichia coli*. *Gene* 144, (1994), 69-73.

Allen, P., Worland, S. and Gold, L.: Isolation of High-Affinity RNA Ligands to HIV-1 Integrase from a Random Pool. *Virology* 209 (1995), 327-336.

Basler, K., Oesch, B., Scott, M., Westaway, D., Wälchli, M., Groth, D.F., McKinley, M.P., Prusiner, S.B. and Weissmann, C.: Scrapie and cellular PrP isoforms are encoded by the same chromosomal gene. *Cell* 46 (1986), 417-428.

Bock, L.C. et al., Selection of single-stranded DNA molecules that bind and inhibit human thrombin. *Nature* 355 (1992), 564-566.

Brandner, S. et al., Normal host prion protein necessary for scrapie-induced neurotoxicity. *Nature* 379 (1996), 339-343.

Büeler, H. et al., Mice devoid of PrP are resistant to scrapie. *Cell* 73 (1993), 1339-1347

Carr, D.W., Stofko-Hahn, R.E., Fraser, I.D.C., Bishop, S.M., Acott, T.S., Brennan, R.G. and Scott, J.D., Interaction of the regulatory subunit (RII) of cAMP-dependent protein kinase with RII-anchoring proteins occurs through an amphipathic helix binding motif. *J. Biol. Chem.* 266 (1991), 14188-14192.

Caughey, B. et al., N-terminal truncation of the Scrapie-associated form of PrP by lysosomal protease(s): Implications regarding the site of conversion of PrP to the protease-resistant state. *J. Virol.* 65 (1991), 6597-6603.

Collinge, J. et al., Prion protein is necessary for normal synaptic function. *Nature* 370 (1994), 295-297.

Edenhofer, F. et al., Prion protein PrP^c interacts with molecular chaperones of the Hsp60 family. *J. Virol.* 70 (1996), 4724-4728.

Epstein, L., Cai, J., Glaser, T., Jepeal, L. and Maas, R.: Identification of a *Pax* paired Domain recognition sequence and evidence for DNA-dependent conformational changes. *J. Biol. Chem.* 269 (1994), 8355-8361.

Famulok, M. and Szostak, J.W.: In vitro Selection of specific Ligand-binding Nucleic Acids. *Angew. Chem. Int. Ed. Engl.* 31 (1992), 979-988.

Famulok, M.: Molecular Recognition of Amino Acids by RNA-Aptamers: An L-Citrulline Binding Motif and its Evolution into an L-Arginine Binder. *J. Am. Chem. Soc.* 116 (1994), 1698-1706.

Groschup, M.H., Langeveld, J. and Pfaff, E.: The major species specific epitope in prion proteins of ruminants. *Arch. Virol.* 136 (1994), 423-431.

Hampton, R.Y., Koning, A., Wright, R., and Rine, J., In vivo examination of membrane protein localization and degradation with green-fluorescent protein. *Proc. Natl. Acad. Sci. USA* 93 (1996), 828-833.

Harada, K. and Frankel, A.D., Identification of two novel arginine binding DNAs. *EMBO J.* 14 (1995), 5798-5811

Hope, J., Morton, L.J.D., Farquhar, C.F., Multhaup, G., Beyreuther, K. and Kimberlin, R.H., The major polypeptide of scrapie-associated fibrils (SAF) has the same size, charge distribution and N-terminal protein sequence as predicted for the normal brain protein (PrP). *EMBO J.* 5 (1986), 2591-2597.

Huizenga, D.E. and Szostak, J.W., A DNA aptamer that binds adenosine and ATP. *Biochemistry* 34 (1995), 656-665

Kawase, M., Mornoeda, M., Young, N.S. and Kijagaya, S., Most of the VP1 unique region of B19 parvovirus is on the capsid surface. *Virology* 211 (1995), 359-366.

Kellings, K., Meyer, N., Mirenda, C., Prusiner, S.B., and Riesner, D.: Further analysis of nucleic acids in purified scrapie prion preparations by improved return refocusing gel electrophoresis. *J. Gen. Virol.* 73 (1992), 1025-1029.

Kim, J.S. and Raines, R.T., Ribonuclease S-peptide as a carrier in fusion proteins. *Proteins Sci.* 2 (1993), 348-356.

Kubik, M.F., Stephens, A.W., Schneider, D., Marlar, R.A. and Tasset, D.: High-affinity RNA ligands to human α -thrombin. *Nucleic Acids Res.* 22 (1994), 2619-2626.

Lansbury, P.T.Jr. and Caughey, B.: The chemistry of scrapie infection: implications of the 'ice 9' metaphor. *Chemistry & Biology* 2 (1995), 1-5 .

Lauhon, C.T. and Szostak, J.W., RNA aptamers that bind flavin and nicotinamide redox cofactors. *J. Am. Chem. Soc.* 117 (1995), 1246-1257

Le-Grice, S.F. and Gruninger-Leitch, F., Rapid purification of homodimer and heterodimer HIV-1 reverse transcriptase by metal chelate affinity chromatography. *Eur. J. Biochem.* 187 (1990), 307-314.

Macaya, R.F. et al., Thrombin-binding DNA aptamer forms a unimolecular quadruplex structure in solution. *Proc. Natl. Acad. Sci. USA* 90 (1993), 3745-3749.

Meisterernst, M., Gander, I., Rogge, L. and Winnacker, E.-L.: A quantitative analysis of nuclear factor I/DNA interactions. *Nucleic acids Res.* 16 (1988), 4419-4435.

Oesch, B., Westaway, D., Wälchli, M., McKinley, M.P., Kent, S.B., Aebersold, R., Barry, R.A., Tempst, B., Teplow, D.B., Hood, L.E., Prusiner, S.B. and Weissmann, C.: A cellular gene encodes scrapie PrP²⁷⁻³⁰ protein. *Cell* 40 (1985), 735-746.

Pan, K.-M., Baldwin, M., Nguyen, J., Gasset, M., Serban, A., Groth, D., Mehlhorn, I., Huang, Z., Fletterick, R.J., Cohen F.E. and Prusiner, S.B.: Conversion of α -helices into β -sheets features in the formation of the scrapie prion proteins. *Proc. Natl. Acad. Sci. USA* 90 (1993), 10962-10966.

Prusiner, S.B., McKinley, M.P., Groth, D.F., Bowman, K.A., Mack, N.I., Cochran, S.P. and Masiarz, F.R.: Scrapie agent contains a hydrophobic protein. *Proc. Natl. Acad. Sci. USA* 78 (1981), 6675-6679.

Prusiner, S.B.: Novel proteinaceous infectious particles cause scrapie. *Science* 216 (1982), 136-144.

Prusiner, S.B., McKinley, M.P., Bowman, K.A., Bolton, D.C., Bendheim, P.E., Groth, D.F. and Glenner, G.G.: Scrapie prions aggregate to form amyloid-like birefringent rods. *Cell* 35 (1983), 349-358.

Prusiner, S.B., Groth, D.F., Bolton, D.C., Kent, S.B., and Hood, L.E.: Purification and structural studies of a major scrapie prion protein. *Cell* 38 (1984), 127-134.

Richards, F.M. and Wyckoff, H.W. in "The Enzymes" Vol IV (Boyer, P.D., Ed.) (1971), 647-806, Academic Press, New York.

Saiki, R.K., Gelfand, D.H., Stoffel, S., Scharf, S.J., Higuchi, R., Horn, G.T., Mullis, K.B. and Erlich, H.A.: Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 239 (1988), 487-491.

Sakaguchi, S. et al., Loss of cerebellar Purkinje cells in aged mice homozygous for disrupted *PrP* gene. *Nature* 380 (1996), 528-531.

Sakashita, E. and Sakamoto, H.: Characterization of RNA binding specificity of the *Drosophila* sex-lethal protein by *in vitro* ligand selection. *Nucleic Acids. Res.* 22 (1994), 4082-4086.

Sambrook J., Fritsch, E.F. and Maniatis, T.: Molecular Cloning. A Laboratory Manual, 2nd ed. Cold Spring Harbor Laboratory Press. Cold Spring Harbor, NY (1989).

Sanger, F., Nicklen, S. and Coulson, A.R.: DNA sequencing with chain terminating inhibitors. *Proc. Natl. Acad. Sci. USA* 74 (1977), 5463-5467.

Schellenberger, A., Fischer, G., Hübner, G. and Ulbrich, R.: Enzymkatalyse. VEB Gustav Fischer Verlag, Jena (1989).

Sen, D. and Gilbert, W., *Nature* 334 (1988), 364-366

Stahl, N., Baldwin, M.A., Teplow, D.B., Hood, L., Gibson, B.W., Burlingame, A.L. and Prusiner, S.B.: Structural studies of the scrapie prion protein using mass spectrometry and amino acid sequencing. *Biochemistry* 32 (1993), 1991-2002.

Sundquist, W.I. and Klug, A., Telomeric DNA dimerizes by formation of guanine tetrads between hairpin-loops. *Nature* 342 (1989), 825-829

Tobler, I. et al., Altered circadian activity rhythms and sleep in mice devoid of prion protein. *Nature* 380 (1996), 639-642

Tuerk, C. and Gold, L.: Systematic Evolution of Ligands by Exponential Enrichment: RNA Ligands to Bacteriophage T4 DNA polymerase. *Science* 249 (1990), 505-510.

Tuerk, C., MacDougall, S. and Gold, L.: RNA pseudoknots that inhibit human immunodeficiency virus type 1 reverse transcriptase. *Proc. Natl. Acad. Sci.* 89 (1992), 6988-6992.

Wang, K.Y. et al., The tertiary structure of a DNA aptamer which binds to and inhibits thrombin determines activity. *Biochemistry* 32 (1993), 11285-11292

Wang, L.-F., Yu, M., White, J.R. and Eaton, B.T., BTag: a novel six-residue epitope tag for surveillance and purification of recombinant proteins. *Gene* 169 (1996), 53-58.

Weiss, S. et al., The multimerization state of retroviral RNA is modulated by ammonium ions and effects HIV-1 full-length cDNA synthesis in vitro. *Nucl. Acids Res.* 21 (1993), 4879-4885.

Weiss, S., König, B., Müller, H.-J., Seidel, H. and Goody, R.S.: Synthetic human tRNA^{Lys3} and natural bovine tRNA^{Lys3} interact with HIV-1 reverse transcriptase and serve as specific primers for retroviral cDNA synthesis. *Gene* 111 (1992), 183-197.

Weiss, S., Famulok, M., Edenhofer, F., Wang, Y.-H., Jones, I.M., Groschup, M., and Winnacker, E.-L.: Overexpression of Active Syrian Golden Hamster Prion Protein PrP^C as a Glutathione S-Transferase Fusion in Heterologous Systems. *J. Virol.* 69 (1995), 4776-4783.

Weiss, S., Rieger, R., Edenhofer, F., Fisch, E. and Winnacker, E.-L.: Recombinant Prion Protein rPrP27-30 from Syrian Golden Hamster reveals Proteinase K Sensitivity, *Biochem. Biophys. Res. Commun.* 219 (1996), 173-179

Williamson, J.R. et al., Monovalent cation-induced structure of telomeric DNA: The G-quartet model. *Cell* 59 (1989), 871-880

Williamson, J.R. et al., G-quartets in biology: A reprise. *Proc. Natl. Acad. Sci. USA* 90 (1993), 3124

Yoshimoto, J., Iinuma, T., Ishiguro, N., Horiuchi, M., Imamura, M. and Shinagawa, M., Comparative sequence analysis and expression of bovine PrP gene in mouse L-929 cells. *Virus genes* 6 (1992), 343-356.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: Winnacker, Ernst-Ludwig
- (B) STREET: Dall Armistrasse 41a
- (C) CITY: Munich
- (E) COUNTRY: Germany
- (F) POSTAL CODE (ZIP): 80638

(ii) TITLE OF INVENTION: Nucleic acid molecules capable of distinguishing the isoforms PrPc and PrPSc of prion proteins and processes for their production

(iii) NUMBER OF SEQUENCES: 20

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 86 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CACUGCAGCA AUUCGUUGUG CGGGAAUUUG AGGGACGAUG GGAAGUGGG GACGAAUGAC 60

UCAUUGCCGC GGUAGGGUUA GGCACC 86

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 86 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

30

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CACUGCAGCA AUGCGUUGUG UGGGAAUUUG AGGGACGAUG GGGAAGUGGG GACGAAUGAC 60
UCAUUGCCGC GGUAGGGUUA GGCACC 86

(2) INFORMATION FOR SEQ ID NO: 3:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 86 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

CACUGC UACC UUAGAGUAGG AGCGGGACGA GGGGUUGUUG GGACGUGGGU AUGAUCCAUA 60
CAUUGAGGAAG CUGGUGAGCU GGCACC 86

(2) INFORMATION FOR SEQ ID NO: 4:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 86 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

CACUGCUACC UUAGAGUAGG AGCGGGACGA GGGGUUGUUG GGACGUGGGU AUGAUCCAUA 60
CAUUAGGAAG CUGGUGAGCU GGCACC 86

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 86 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

CACUGCUACC UUAGAGUAGG AGCGGGACGA GGGGUUGUUG GGACGUGGGU AUGAUCCAUA 60
CAUUAGGAAG CUGGUGAGCU GGCACC 86

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 81 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

CACUGCGACA UGGGAAGAGG GAAGAGGGUU GUCGGGAGAU AAUGUCGCGA AACUAAGAAC 60
UCUAAGAGCU GCCCGGGCAC C 81

32

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 82 bas pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

CACUGCGACA UGGGAGGAGG GAAGAGGGUU GUCGGGAGAU AAUGUCGCGA AACUAAGAAC	60
UCUAAGAGCU GCCCGUGGCA CC	82

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 83 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

CACUGCGACA UGGGAAGAGG GAAGAGGGUU GUCGGGAGAU AAUGUCGCAA AGCUAAGAAC	60
UCUAAGAGCU GCCGCGUGGC ACC	83

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 85 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /d sc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

CACUGCUUGC UCGUUGCACU GUGAUAUGUG GGUUUAGGAU AGGGAGAAGG GAAGAGGGAA 60

GAAUAUCCGU CUGAACGAGG GCACC 85

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 85 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

CACUGCUGCU AUUCAGUGGG UUGUGGGAGA AGGUAGGGG GAUGAUGAAA GCAGCUCGUG 60

UGAUUUCUUU CUGAAGACCG GCACC 85

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 85 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

CACUGCCGUC AUAUGGGCAC AUCUCAAGU GGGAAUGUGG GGUGAUGGGA AGAGGGAUGA 60

UUAAGAUGGC CACAUAUUCG GCACC 85

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 85 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

CACUGCGAGG AUGCGGGACG AGGGAACGUG GGAACGAGGG AUGAAUCCUU GUAGUGAGAU	60
AGCUUCCCCA ACAUGUCCAG GCACC	85

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 85 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

CACUGCUCGC GUCAUUGGCA AGAGGGAAGU GGAUGCGGG AAAGAUUGGG AACACCGCAC	60
CAAUAAUGUG AGUGUGAGGG GCACC	85

(2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 81 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

CACUGCCUCG AAAACUGUGA AGAGUACGCU UUAACUGUGC UCCGUGUGGA UUGACCAUAG 60

ACCCGUCCCU GGACAGGCAC C 81

(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

GGGAUUUGA GGGACGAUGG GGAAGUGGG 29

(2) INFORMATION FOR SEQ ID NO: 16:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

GGGACGAGGG GUUGUUGGGA CGUGGG 26

36

(2) INFORMATION FOR SEQ ID NO: 17:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

GGGAAGAGGG AAGAGGGUUG UCGGG

25

(2) INFORMATION FOR SEQ ID NO: 18:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 60 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "RNA"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

GCA AUGCGUU GUGUGGGAU UUGAGGGACG AUGGGGAAGU GGGGACGAU GACUCAUUGC

60

(2) INFORMATION FOR SEQ ID NO: 19:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 44 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "oligodesoxynucleotide"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

37

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

CCGAATTCTA ATACGACTCA CTATAGGAGC TCAGCCTTCA CTGC

44

(2) INFORMATION FOR SEQ ID NO: 20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligodesoxynucleotide"

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

GTGGATCCGA CCGTGGTGCC

20

Claims

1. A process for the identification and isolation of nucleic acid molecules which are capable of distinguishing between the isoforms PrP^c and PrP^{Sc} of prion proteins associated with transmissible spongiform encephalopathies comprising the steps of
 - (i) incubating a prion protein isoform or peptide fragment or derivative of this prion protein isoform with a pool of nucleic acid molecules comprising different sequences;
 - (ii) selecting and isolating those nucleic acid molecules which are capable of binding to said prion protein isoform or fragment or derivative thereof;
 - (iii) optionally, amplifying the isolated nucleic acid molecules and repeating steps (i) and (ii); and
 - (iv) determining the binding specificity of the isolated nucleic acid molecules for the PrP^c and PrP^{Sc} isoforms of prion proteins.
2. The process according to claim 1, wherein the nucleic acid molecule is RNA.
3. The process according to claim 1 or 2, wherein the pool of nucleic acid molecules is a randomized RNA pool.
4. The process according to claim 3, wherein the pool of nucleic acid molecules is the RNA pool M 111.1 as described in Famulok (J. Am. Chem. Soc. 116 (1994), 1698-1706).

5. The process according to claim 1, wherein the nucleic acid molecule is DNA.
6. The process according to claim 5, wherein the DNA is single stranded DNA.
7. The process according to claim 5, wherein the DNA is double stranded DNA.
8. The process according to any one of claims 5 to 7, wherein the pool of nucleic acid molecules is a randomized DNA pool.
9. The process according to any one of claims 1 to 8 wherein the transmissible spongiform encephalopathy is Scrapie, bovine spongiform encephalopathy (BSE), Creutzfeld-Jacob Disease (CJD), Gerstmann-Sträußler-Scheinker-Syndrome (GSS), Kuru, fatal familial insomnia (FFI) or transmissible mink encephalopathy (TME), feline spongiform encephalopathy (FSE) or chronic wasting disease (CWD).
10. The process according to any one of claims 1 to 9, wherein the prion protein isoform or fragment or derivative thereof is immobilized.
11. The process according to any one of claims 1 to 9, wherein the prion protein isoform or fragment or derivative thereof is in solution.
12. The process according to any one of claims 1 to 11, wherein the prion protein isoform is the isoform PrP^{Sc} or a fragment or derivative of this isoform.

13. The process according to claim 12, wherein the prion protein is the derivative PrP²⁷⁻³⁰ or a fragment thereof.
14. The process according to any one of claims 1 to 11, wherein the prion protein isoform is the isoform PrP^c or a fragment or derivative of this isoform.
15. The process according to claim 14, wherein the prion protein is the processed form PrP^c₂₃₋₂₃₁.
16. The process according to any one of claims 1 to 15, wherein the prion protein is a recombinant protein.
17. The process according to claim 16, wherein the prion protein is part of a fusion protein.
18. The process according to claim 17, wherein the prion protein is part of a fusion protein with oligohistidine, calmoduline binding protein, S-peptide, FLAG, green-fluorescent protein, BTag, maltose binding protein or glutathione-S-transferase.
19. A nucleic acid molecule obtainable by the process according to any one of claims 1 to 18.
20. The nucleic acid molecule according to claim 19 which specifically binds to an isoform of a prion protein as defined in any one of claims 10 to 15 or to a fragment or derivative thereof.

21. The nucleic acid molecule according to claim 19 or 20 which is an RNA molecule.
22. The nucleic acid molecule according to claim 19 or 20 which is a DNA molecule.
23. The nucleic acid molecule according to any one of claims 19 to 22 which comprises four stretches of three consecutive guanosine residues separated by regions between four and seven nucleotides long.
24. The nucleic acid molecule according to claim 23 which comprises a nucleotide sequence as depicted in SEQ ID NO: 15, SEQ ID NO: 16, or SEQ ID NO: 17.
25. The nucleic acid molecule according to any one of claims 19 to 23, which comprises a nucleotide sequence as depicted in any one of SEQ ID NO: 1 to SEQ ID NO: 13.
26. The nucleic acid molecule according to any one of claims 19 to 22 which comprises a nucleotide sequence as depicted in SEQ ID NO: 18.
27. The nucleic acid molecule according to any one of claims 19 to 26 which is modified at one or more positions in order to increase its stability and/or to alter its biophysical and/or biochemical properties.
28. A pharmaceutical composition comprising a nucleic acid molecule according to any one of claims 19 to 27 and optionally a pharmaceutically acceptable carrier.

29. A diagnostic composition comprising a nucleic acid molecule according to any one of claims 19 to 27.
30. A method for the in-vitro diagnosis of a transmissible spongiform encephalopathy, wherein at least one of the nucleic acid molecules according to any one of claims 19 to 27 is incubated with a probe and the interaction between the nucleic acid molecules and the PrP^c or PrP^{Sc} isoforms of a prion protein or fragment or derivative of these isoforms is determined.
31. The method according to claim 30 wherein at least one of the nucleic acid molecules according to any one of claims 19 to 27 is used to quantitatively determine the amount of at least one isoform of a prion protein or a fragment or derivative thereof in a probe.
32. The method according to claim 31, wherein nucleic acid molecules are used which specifically bind to the cellular isoform PrP^c or fragment or derivatives thereof in combination with nucleic acid molecules which specifically bind to the isoform PrP^{Sc} or fragment or derivatives thereof and the absolute and/or relative amount of the isoforms PrP^c and PrP^{Sc} in the probe is determined.
33. The method according to any one of claims 30 to 32, wherein the probe is from an organ tissue.
34. The method of claim 32 wherein the organ tissue is from brain, tonsils, ileum, cortex, dura mater, Purkinje cells, lymphnodes, nerve cells, spleen, muscle cells, placenta, pancreas, eyes, backbone marrow or peyer'sche plaqu .

35. The method according to any one of claims 30 to 32, wherein the probe is from a body fluid.
36. The method according to claim 35, wherein the body fluid is blood, cerebrospinal fluid, milk or semen.
37. A chemical compound other than a nucleic acid molecule based on information derived from a three dimensional structure of nucleic acid molecules according to any one of claims 19 to 27 selected from the group consisting of inorganic or organic compounds.
38. The chemical compound according to claim 37, which is a sugar, an amino acid, a protein or a carbohydrate.

1/11




Fig. 1 A

2/11







Fig. 1 B

3/11

**Fig. 2**

4/11

**Fig. 3**

**Fig. 4 A**


6/11

a

selected RNA (after 9 Cycles)

GST GST:: GST::
rPrP^c rPrP27-30

1:1 10:1 1:1 10:1 1:1 10:1



-RNA/
GST::rPrP^c
Interaction

-free RNA

1 2 3 4 5 6 7 8 9

b

selected RNA (after 11 Cycles)

GST GST:: GST::
rPrP^c rPrP27-30

1:1 10:1 1:1 10:1 1:1 10:1



molar ratio
RNA : protein

-RNA/
GST::rPrP^c
Interaction

-free RNA

1 2 3 4 5 6 7 8 9

Fig. 4 B

7/11

A

#I CACUGCAGCAAUUCGUUGUGCGGGAUUUGAGGGACGAU~~GGGA~~AGUGGGGACGAAUGACUCA
UUGCCGCGGUAGGGUJAGGCACC

#II CACUGCAGCAAUUGCGUUGUGUGGGAAUUUGAGGGACGAU~~GGGA~~AGUGGGGACGAAUGACUC
AUUGCCGCGGUAGGGUJAGGCACC

B

#III CACUGCUACCUUAGAGUAGGAGCGGGACGAGGGGUUGUU~~GGGA~~CGUGGGUAUGAUCCAUAUAC
AUUAGGAAGCUGGUGAGCUGGCACC

#IV CACUGCUACCUUAGAGUAGGAGCGGGACGAGGGGUUGUU~~GGGA~~CGUGGGUAUGAUCCAUAUAC
AUUAGGAAGCUGGUGAGCUGGCACC

#V CACUGCUACCUUAGAGUAGGAGCGGGACGAGGGGUUGUU~~GGGA~~CGUGGGUAUGAUCCAUAUAC
AUUAGGAAGCUGGUGAGCUGGCACC

C

#VI CACUGCGACAUGGGGAAGAGGGGUUGUCGGGAGAUAAUGUCGCGAAACUAAGAACU
CUAAGAGCUGCCCGTGGCACC

#VII CACUGCGACAUGGGGAGGAGGGGUUGUCGGGAGAUAAUGUCGCGAAACUAAGAAC
UCUAAGAGCUGCCCGUGGCACC

#VIII CACUGCGACAUGGGGAAGAGGGGUUGUCGGGAGAUAAUGUCGCGAAAGCUAAGAAC
UCUAAGAGCUGCCCGUGGCACC

fig 5

D

- IX CACUGCUUGCUCGUGCACUGUGAU AUGUGGGUUAAGGAUAGGGAGAGGGGAAGAGGGGAAGAAUAUCCGUCU
GAACGAGGGCACC
- X CACUGCUGCUAUUCAGUGGGUUGUGGGAGAGGGUAGGGGAUGAUGAAAGCAGCUCGUGUGAUUUCUUCUGAA
GACCGGCACC
- XI CACUGCCGUCAU AUGGGCACAUUCUCAAAGUGGGAAUGUGGGGUGAUGGGAAGAGGGAUGAUUAAGAUGGGCCACAUA
UUCGGCACC
- XII CACUGCGAGGAUGCGGGACGAGGGAACGUGGGGAUGAAUCCUUGUAGUGAGAUAGCUUCCCCCAACAUG
UCCAGGCACC
- XIII CACUGCUCGCGUCAUUGGCAAGAGGGGAAGUGGGGAUGCGGGAAAGAUUGGGAACACCGCACCAUAUAGUGAGUGU
GAGGGGCACC

E

- XIV CACUGCCUCGAAACUGUGAGAGUACGCUUUAACUGUGCUCGUGUGGAUUGACCAUAGACCCGUCGCCUGG
ACAGGCACC

fig 5

9/11






Fig. 6



A Aptamer Motif I



B Aptamer Motif II



Fig. 7




Fig. 7

INTERNATIONAL SEARCH REPORT

International Application No

PC/EP 96/04671

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12Q1/68 C07H21/02 C07H21/04 A61K31/70 G01N33/53

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12Q C07H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	VETINARY QUARTERLY, vol. 15, no. 4, December 1993, pages 167-74, XP000617784 SCHREUDER B. E.: "General aspects of transmissible spongiform encephalopathies and hypotheses about the agents" see the whole document ---	1
A	WO,A,93 20242 (THE SCRIPPS RESEARCH INSTITUTE) 14 October 1993 see claims 1-19 --- -/--	1-8, 18-29, 37,38

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

A document member of the same patent family

2

Date of the actual completion of the international search 14 February 1997	Date of mailing of the international search report 18.03.97
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer Osborne, H

INTERNATIONAL SEARCH REPORT

International Application No

PC1/EP 96/04671

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>MOLECULAR BIOLOGY REPORTS, vol. 20, no. 2, 1994, pages 97-107, XP000617781 KLUG S ET AL: "All you wanted to know about SELEX" see the whole document ---</p>	1-8, 18-29
A	<p>THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 23, June 1995, pages 13581-84, XP002025263 GOLD L: "Oligonucleotides as research, diagnostic, and therapeutic agents " see the whole document ---</p>	1-8, 18-29
A	<p>JOURNAL OF VIROLOGY, vol. 69, no. 8, August 1995, pages 4776-83, XP000644456 WEISS S ET AL: "Over expression of active golden hamster prion protein PrPc as a glutathionine S-transferase fusion in heterologous systems" cited in the application see the whole document ---</p>	1,16-18
A	<p>J. AM. CHEM. SOC., vol. 116, March 1994, pages 1698-1706, XP000644374 FAMULOK M: "Molecular recognition of amino acids by RNA-Aptamers: An L-citrulline binding RNA motif and its evolution into an L-arginine binder" cited in the application see the whole document -----</p>	1-8, 18-29

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC 1/EP 96/04671

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9320242	14-10-93	US-A- 5573905	12-11-96
		AU-A- 3944993	08-11-93
		CA-A- 2132103	14-10-93
		EP-A- 0643778	22-03-95
		JP-T- 7505530	22-06-95
